



rec'd from
Dorset Research
Centre
Apr 15 86

05/15/86
TWB

MORPHOMETRY
OF THE MUSKOKA-HALIBURTON
STUDY LAKES

A. Nicolls, R. Reid and **R.** Girard

DATA REPORT DR 83/3

TD
227
.H35
N53
1983
MOE

Copyright Provisions and Restrictions on Copying:

This Ontario Ministry of the Environment work is protected by Crown copyright (unless otherwise indicated), which is held by the Queen's Printer for Ontario. It may be reproduced for non-commercial purposes if credit is given and Crown copyright is acknowledged.

It may not be reproduced, in all or in part, for any commercial purpose except under a licence from the Queen's Printer for Ontario.

For information on reproducing Government of Ontario works, please contact Service Ontario Publications at copyright@ontario.ca

DATA REPORT SERIES

The data presented in this report were collected by staff of the Water Resources Branch of the Ontario Ministry of the Environment as part of the Lakeshore Capacity Study or the Acid Precipitation in Ontario Study. This unreviewed report does not necessarily reflect the views or opinions of the Ontario Ministry of the Environment.

MORPHOMETRY OF THE MUSKOKA-HALIBURTON
STUDY LAKES

A. Nicolls, R. Reid and R. Girard
Limnology Unit,
Water Resources Branch,
Ontario Ministry of the Environment,
P.O. Box 39, DORSET, Ontario
POA 1E0


DATA REPORT DR 83/3

PREFACE

The unpublished Data Report Series is intended as a readily available source of basic data collected for lakes and watersheds in the Muskoka-Haliburton area of Ontario. These data were collected as part of the Lakeshore Capacity Study and/or the Acid Precipitation in Ontario Study.

The limnological portion of the Lakeshore Capacity Study (1975-81) was initiated to investigate the relationships between lakeshore development and lake trophic status in low ionic strength Precambrian lakes. The Acid Precipitation in Ontario Study (1979-present) was initiated, in part, to investigate the effects of the deposition of strong acids on aquatic and terrestrial ecosystems in Ontario. The primary findings of these studies have been and will continue to be published as reviewed papers and technical reports.

ABSTRACT

 Morphometric maps and tables are presented for the lakes studied as part of the Lakeshore Capacity Study.

Nicolls, A., R. Reid and R. Girard. 1983. Morphometry of the Muskoka-Haliburton Study Lakes. Ont. Min. Envir. Data Report DR 83/3.

TABLE OF CONTENTS

	<u>Page</u>
INTRODUCTION.....	1
BLUE CHALK LAKE.....	4
CHUB LAKE.....	6
DICKIE LAKE.....	8
HARP LAKE.....	10
JERRY LAKE.....	12
RED CHALK LAKE.....	14
BASSHAUNT LAKE.....	16
BIGWIND LAKE.....	18
BUCK LAKE.....	20
CROSSON LAKE.....	22
GLEN LAKE.....	24
GULLFEATHER LAKE.....	26
LITTLE CLEAR LAKE.....	28
SOLITAIRE LAKE.....	30
WALKER LAKE.....	32

Introduction

This report summarizes the morphometric information collected for the study lakes of the trophic status component of the Lakeshore Capacity Study. The methodology used to prepare the maps and calculate morphometric parameters is outlined by Scheider et al. (1983). The lake outlines were taken from aerial photographs; more accurate ground surveys will be carried out in 1983-84 and may result in slight changes in lake areas and volumes.

The morphometric parameters calculated for each lake are defined by Hutchinson (1957) and include the following:

- | | |
|---------------------------------------|---|
| 1) Area (A) | - Lake surface area. |
| 2) Volume (V) | - Total lake volume calculated by summing the individual stratum volumes determined as noted below. |
| 3) Mean Depth (\bar{z}) | - Calculated as V/A . |
| 4) Maximum Depth (z_m) | |
| 5) Shoreline Length (L) | - Total shoreline length. |
| 6) Development of Shoreline (D_L) | - The ratio (dimensionless) of the shoreline length to the length of the circumference of a circle of equal area to that of the lake: |

$$D_L = L/2 (\pi A)^{1/2}$$

D_L must be >1 and is a measure of the irregularity of the shoreline.

- | | |
|------------------------------------|--|
| 7) Development of Volume (D_V) | - The ratio of the volume of the lake to that of a cone of basal area A and height z_m : |
|------------------------------------|--|

$$D_V = 3\bar{z}/z_m$$

This quantity is an expression of the form of the lake basin.

- 8) Contour Area (A_i) - The surface area within each of the depth contours. These contour areas are used to calculate strata volumes.
- 9) Stratum Volume - The volume of a lake stratum between two contour depths:

$$V_{m-n} = 1/3 (A_m + A_n + (A_m A_n)^{1/2}) (m-n)$$

where m = the lower contour depth, and
n = the upper contour depth.

A morphometric map and a table of morphometric information follow for each lake. Since Red Chalk Lake can be divided into two distinct basins, the morphometry for each basin has also been provided.

References

- Hutchinson, G.E. 1957. A Treatise on Limnology. John Wiley and Sons, Inc., 1015 p.
- Scheider, W.A., R.A. Reid, B. Locke and L.D. Scott. 1983. Studies of lakes and watersheds in Muskoka-Haliburton, Ontario: Methodology. Ont. Min. Envir. Data Report 83/1.

Inflow 1

BLUE CHALK LAKE

MUSKOKA Dist.

RIDOUT Tp.

Lat. 45° 12' Long. 78° 56'



0m 100m

500m

Table 1:

BLUE CHALK LAKE MORPHOMETRY SUMMARY

Lake Area A (ha)	Lake Volume V (m ³ x10 ⁵)	Mean Depth \bar{z} (m)	Maximum Depth Z _m (m)	Shoreline Length L (km)	Development of Shoreline D _L	Development of Volume D _V
49.4	42.1	8.5	23	4.60	1.85	1.11

Contour Depth (m)	Contour Area (ha)	Stratum Volume (m ³ x10 ⁵)
0	49.4	8.89
2	39.7	7.38
4	34.2	6.35
6	29.4	5.24
8	23.1	3.95
10	16.6	3.10
12	14.5	2.63
14	11.9	2.15
16	9.64	1.41
18	4.73	0.741
20	2.76	0.275
22	0.359	0.012
23	0	




Table 2:

CHUB LAKE MORPHOMETRY SUMMARY

Lake Area A (ha)	Lake Volume V ($m^3 \times 10^5$)	Mean Depth \bar{z} (m)	Maximum Depth Zm (m)	Shoreline Length L (km)	Development of Shoreline D_L	Development of Volume D_V
32.2	28.5	8.9	27	3.87	1.92	0.99

Contour Depth (m)	Contour Area (ha)	Stratum Volume ($m^3 \times 10^5$)
0	32.2	5.77
2	25.6	4.73
4	21.7	3.91
6	17.5	3.17
8	14.3	2.64
10	12.1	2.23
12	10.2	1.78
14	7.72	1.37
16	5.98	1.06
18	4.61	0.782
20	3.25	0.558
22	2.36	0.355
24	1.25	0.115
26	0.109	0.004
27	0	





Table 3:

DICKIE LAKE MORPHOMETRY SUMMARY

Lake Area A (ha)	Lake Volume V (m ³ ×10 ⁵)	Mean Depth \bar{z} (m)	Maximum Depth Zm (m)	Shoreline Length L (km)	Development of Shoreline D _L	Development of Volume D _V
93.2	46.4	5.0	12	7.84	2.29	1.25

Contour Depth (m)	Contour Area (ha)	Stratum Volume (m ³ ×10 ⁵)
0	93.2	16.7
2	74.5	6.99
3	65.4	6.10
4	56.7	4.84
5	40.6	3.73
6	34.1	3.04
7	26.9	2.27
8	18.7	1.55
9	12.5	0.883
10	5.58	0.280
11	0.752	0.025
12	0	



HARP LAKE

MUSKOKA Dist.

CHAFFEY Tp.

Lat. 45° 23' Long. 79° 08'



Table 4:

HARP LAKE MORPHOMETRY SUMMARY

Lake Area A (ha)	Lake Volume V (m ³ x10 ⁵)	Mean Depth \bar{Z} (m)	Maximum Depth Z _m (m)	Shoreline Length L (km)	Development of Shoreline D _L	Development of Volume D _V
66.9	82.6	12.4	40	4.56	1.57	0.93

Contour Depth (m)	Contour Area (ha)	Stratum Volume (m ³ x10 ⁵)
0	66.9	12.3
2	56.5	10.7
4	50.3	9.4
6	43.7	8.32
8	39.5	7.37
10	34.2	6.35
12	29.4	5.46
14	25.3	4.73
16	22.0	4.10
18	19.1	3.51
20	16.1	2.90
22	13.0	2.33
24	10.3	1.79
26	7.7	1.28
28	5.17	0.895
30	3.81	0.559
32	1.89	0.313
34	1.27	0.194
36	0.703	0.092
38	0.250	0.017
40	0	




Table 5:

JERRY LAKE MORPHOMETRY SUMMARY

Lake Area A (ha)	Lake Volume V (m ³ x10 ⁵)	Mean Depth \bar{Z} (m)	Maximum Depth Zm (m)	Shoreline Length L (km)	Development of Shoreline D _L	Development of Volume D _V
50.1	61.9	12.4	35	4.60	1.83	1.06

Contour Depth (m)	Contour Area (ha)	Stratum Volume (m ³ x10 ⁵)
0	50.1	9.51
2	45.0	8.54
4	40.5	14.5
8	32.3	11.4
12	24.8	8.39
16	17.3	3.13
18	14.0	2.41
20	10.2	2.57
24	3.28	0.955
28	1.59	0.433
32	0.64	0.070
34	0.125	0.004
35	0	




Table 6:

Red Chalk Lake Morphometry Summary

	Lake Area A (ha)	Lake Volume V ($m^3 \times 10^5$)	Mean Depth Z (m)	Maximum Depth Zm (m)	Shoreline Length L (km)	Development of Shoreline D_L	Development of Volume D_V
Whole Lake	56.9	80.8	14.2	38	4.82	1.80	1.12
Main Basin	43.9	73.3	16.7	38	3.15	1.34	1.32
East Basin	13.0	7.46	5.7	19	1.67	1.31	0.90

Whole Lake			Main Basin			East Basin		
Contour Depth (m)	Contour Area (ha)	Stratum Volume ($m^3 \times 10^5$)	Contour Depth (m)	Contour Area (ha)	Stratum Volume ($m^3 \times 10^5$)	Contour Depth (m)	Contour Area (ha)	Stratum Volume ($m^3 \times 10^5$)
0	56.9	10.6	0	43.9	8.38	0	13.0	2.17
2	48.7	9.34	2	39.8	7.73	2	8.85	1.61
4	44.8	16.2	4	37.5	14.0	4	7.25	1.28
8	36.7	13.1	8	32.6	12.1	6	5.58	0.965
12	29.0	10.5	12	27.7	10.2	8	4.11	0.653
16	24.0	8.44	16	23.5	8.36	10	2.49	0.372
20	18.4	6.43	20	18.4	6.43	12	1.29	0.202
24	13.8	3.80	24	13.8	3.80	14	0.756	0.125
28	5.75	1.72	28	5.75	1.72	16	0.504	0.074
32	3.01	0.573	32	3.01	0.573	18	0.252	0.008
36	0.315	0.021	36	0.315	0.021	19	0	
38	0		38	0				

BASSHAUNT LAKE

HALIBURTON Co.

GUILFORD Tp.

Lat. 45°07' Long. 78°28'




Table 7:

Basshaunt Lake Morphometry Summary

Lake Area A (ha)	Lake Volume V ($m^3 \times 10^5$)	Mean Depth \bar{z} (m)	Maximum Depth Zm (m)	Shoreline Length L (m)	Development of Shoreline D _L	Development of Volume D _V
47.3	36.6	7.7	24	4.85	1.99	0.96

Contour Depth (m)	Contour Area (ha)	Stratum Volume ($m^3 \times 10^5$)
0	47.3	8.64
2	39.2	7.11
4	32.0	5.57
6	23.9	4.25
8	18.7	3.35
10	14.9	2.63
12	11.5	1.95
14	8.07	1.26
16	4.66	0.824
18	3.60	0.594
20	2.38	0.328
22	1.00	0.091
24	0	

BIGWIND LAKE

MUSKOKA Dist.

OAKLEY Tp.

Lat. $45^{\circ} 03'$ Long. $79^{\circ} 03'$

0m. 500m. 1000m.

N

Outflow




Table 8:

BIGWIND LAKE MORPHOMETRY SUMMARY

Lake Area A (ha)	Lake Volume V (m ³ x10 ⁵)	Mean Depth \bar{z} (m)	Maximum Depth Zm (m)	Shoreline Length L (km)	Development of Shoreline D _L	Development of Volume D _V
111	118	10.7	32	8.24	2.21	1.00

Contour Depth (m)	Contour Area (ha)	Stratum Volume (m ³ x10 ⁵)
0	111	20.8
2	97.2	18.4
4	87.0	28.4
8	56.0	18.3
12	36.3	12.3
16	25.6	8.70
20	18.2	6.09
24	12.5	3.82
28	6.90	1.03
30	3.61	0.241
32	0	

BUCK LAKE

MUSKOKA Dist.

SINCLAIR Tp

Lat. 45° 23' Long 79° 00'




Table 9:

BUCK LAKE MORPHOMETRY SUMMARY

Lake Area A (ha)	Lake Volume V (m ³ x10 ⁵)	Mean Depth \bar{z} (m)	Maximum Depth Zm (m)	Shoreline Length L (km)	Development of Shoreline D _L	Development of Volume D _V
40.3	43.9	10.9	30	3.56	1.58	1.09

Contour Depth (m)	Contour Area (ha)	Stratum Volume (m ³ x10 ⁵)
0	40.3	7.52
2	34.9	6.58
4	31.0	5.77
6	26.8	4.95
8	22.8	4.22
10	19.5	3.63
12	16.8	3.13
14	14.5	2.55
16	11.1	1.90
18	7.95	1.39
20	6.03	0.974
22	3.79	0.596
24	2.24	0.377
26	1.55	0.238
28	0.864	0.083
30	0	

CROSSON LAKE

MUSKOKA Dist.

OAKLEY Tp.

Long. 45° 05' Lat. 79° 02'




Table 10:

CROSSON LAKE MORPHOMETRY SUMMARY

Lake Area A (ha)	Lake Volume V (m ³ x10 ⁵)	Mean Depth \bar{z} (m)	Maximum Depth Zm (m)	Shoreline Length L (km)	Development of Shoreline D _L	Development of Volume D _V
56.8	47.7	8.4	25	3.88	1.45	1.01

Contour Depth (m)	Contour Area (ha)	Stratum Volume (m ³ x10 ⁵)
0	56.8	10.4
2	47.5	8.67
4	39.3	7.07
6	31.5	5.60
8	24.6	4.52
10	20.6	3.66
12	16.1	2.79
14	11.9	2.05
16	8.70	1.44
18	5.84	0.90
20	3.28	0.419
22	1.10	0.132
24	0.304	0.010
25	0	

GLEN LAKE

HALIBURTON Co.

GUILFORD Tp.

Lat. $45^{\circ}08'$ Long. $78^{\circ}30'$




Table 11:

GLEN LAKE MORPHOMETRY SUMMARY

Lake Area A (ha)	Lake Volume V ($m^3 \times 10^5$)	Mean Depth \bar{z} (m)	Maximum Depth Zm (m)	Shoreline Length L (km)	Development of Shoreline D_L	Development of Volume D_V
16.3	11.8	7.2	15	1.83	1.28	1.44

Contour Depth (m)	Contour Area (ha)	Stratum Volume ($m^3 \times 10^5$)
0	16.3	3.02
2	13.9	2.59
4	12.0	2.21
6	10.1	1.79
8	7.81	1.17
10	4.13	0.638
12	2.34	0.321
14	0.976	0.032
15	0	

GULLFEATHER LAKE

MUSKOKA

Dist.

OAKLEY

Tp.

Lat. 45°06' Long. 79°01'




Table 12:

GULLFEATHER LAKE MORPHOMETRY SUMMARY

Lake Area A (ha)	Lake Volume V (m ³ x10 ⁵)	Mean Depth \bar{Z} (m)	Maximum Depth Zm (m)	Shoreline Length L (km)	Development of Shoreline D _L	Development of Volume D _V
65.9	31.5	4.8	13	5.26	1.83	1.11

Contour Depth (m)	Contour Area (ha)	Stratum Volume (m ³ x10 ⁵)
0	65.9	11.6
2	50.8	8.75
4	37.1	5.63
6	20.1	3.25
8	12.6	1.66
10	4.66	0.533
12	1.09	0.036
13	0	

LITTLE CLEAR LAKE

MUSKOKA Dist.

SINCLAIR Tp.

Lat. 45° 24' Long. 79° 00'

TURTLE
LAKE

< 2m.

0m. 100m. 500m.




Table 13:

LITTLE CLEAR LAKE MORPHOMETRY SUMMARY

Lake Area A (ha)	Lake Volume V (m ³ x10 ⁵)	Mean Depth \bar{Z} (m)	Maximum Depth Zm (m)	Shoreline Length L (km)	Development of Shoreline D _L	Development of Volume D _V
10.9	8.86	8.1	25	1.48	1.26	0.97

Contour Depth (m)	Contour Area (ha)	Stratum Volume (m ³ x10 ⁵)
0	10.9	1.99
2	8.99	1.69
4	7.97	1.45
6	6.51	1.19
8	5.39	0.905
10	3.71	0.613
12	2.46	0.395
14	1.52	0.254
16	1.04	0.175
18	0.72	0.116
20	0.448	0.064
22	0.208	0.020
24	0.024	0.001
25	0	

Outflow

SOLITAIRE LAKE

MUSKOKA Dist.

SINCLAIR Tp.

Lat. 45° 22' Long. 79° 00'

Inflow 1




Table 14:

SOLITAIRE LAKE MORPHOMETRY SUMMARY

Lake Area A (ha)	Lake Volume V (m ³ x10 ⁵)	Mean Depth \bar{Z} (m)	Maximum Depth Z _m (m)	Shoreline Length L (km)	Development of Shoreline D _L	Development of Volume D _V
124	164	13.3	31	5.98	1.51	1.29

Contour Depth (m)	Contour Area (ha)	Stratum Volume (m ³ x10 ⁵)
0	124	22.2
2	98.6	18.7
4	88.9	17.1
6	82.3	15.7
8	74.9	14.2
10	66.8	12.7
12	60.6	11.5
14	54.5	10.4
16	49.0	9.27
18	43.7	8.22
20	38.6	7.23
22	33.8	6.24
24	28.7	5.12
26	22.6	3.53
28	13.1	1.56
30	3.52	0.117
31	0	

WALKER LAKE

MUSKOKA Dist.

SINCLAIR Tp

Lat. 45° 24' Long. 79° 05'




Table 15:

WALKER LAKE MORPHOMETRY SUMMARY

Lake Area A (ha)	Lake Volume V (m ³ ×10 ⁵)	Mean Depth \bar{z} (m)	Maximum Depth Z _m (m)	Shoreline Length L (km)	Development of Shoreline D _L	Development of Volume D _V
68.2	42.1	6.2	17	6.44	2.20	1.09

Contour Depth (m)	Contour Area (ha)	Stratum Volume (m ³ ×10 ⁵)
0	68.2	12.6
2	58.2	10.5
4	46.8	8.07
6	34.3	5.32
8	19.6	3.04
10	11.2	1.65
12	5.63	0.715
14	1.86	0.155
16	0.08	0.003
17	0	

**TD
227
.H35
N53
1983**

Morphometry of the Muskoka-
Haliburton study lakes / Nicolls, A.
77942